## Searching For Gravitational Waves From Cosmological Phase Transitions With The NANOGrav 12.5-year dataset

```
Zaven Arzoumanian, Paul T. Baker, Harsha Blumer, 4 Bence Bécsy, Adam Brazier, 6,7 Paul R. Brook, 3,4 Sarah
 Burke-Spolaor, <sup>3, 4, 8</sup> Maria Charisi, <sup>9</sup> Shami Chatterjee, <sup>6</sup> Siyuan Chen, <sup>10, 11, 12</sup> James M. Cordes, <sup>6</sup> Neil J. Cornish, <sup>5</sup>
Fronefield Crawford, <sup>13</sup> H. Thankful Cromartie, <sup>6</sup> Megan E. DeCesar, <sup>14, 15, *</sup> Paul B. Demorest, <sup>16</sup> Timothy Dolch, <sup>17, 18</sup>
 Justin A. Ellis, <sup>19</sup> Elizabeth C. Ferrara, <sup>20, 21, 22</sup> William Fiore, <sup>3, 4</sup> Emmanuel Fonseca, <sup>23</sup> Nathan Garver-Daniels, <sup>3, 4</sup>
 Peter A. Gentile, <sup>3,4</sup> Deborah C. Good, <sup>24</sup> Jeffrey S. Hazboun, <sup>25,*</sup> A. Miguel Holgado, <sup>26,27</sup> Kristina Islo, <sup>28</sup> Ross J.
Jennings, <sup>6</sup> Megan L. Jones, <sup>28</sup> Andrew R. Kaiser, <sup>3,4</sup> David L. Kaplan, <sup>28</sup> Luke Zoltan Kelley, <sup>29</sup> Joey Shapiro Key, <sup>25</sup>
Nima Laal, <sup>30</sup> Michael T. Lam, <sup>31,32</sup> T. Joseph W. Lazio, <sup>33</sup> Vincent S. H. Lee, <sup>34</sup> Duncan R. Lorimer, <sup>3,4</sup> Jing Luo, <sup>35</sup>
     Ryan S. Lynch,<sup>36</sup> Dustin R. Madison,<sup>3,4,*</sup> Maura A. McLaughlin,<sup>3,4</sup> Chiara M. F. Mingarelli,<sup>37,38</sup> Andrea
Mitridate, <sup>34</sup> Cherry Ng, <sup>39</sup> David J. Nice, <sup>14</sup> Timothy T. Pennucci, <sup>40,41,*</sup> Nihan S. Pol, <sup>3,4,9</sup> Scott M. Ransom, <sup>40</sup> Paul
     S. Ray, 42 Brent J. Shapiro-Albert, 3, 4 Xavier Siemens, 30, 28 Joseph Simon, 33, 43 Renée Spiewak, 44 Ingrid H.
   Stairs,<sup>24</sup> Daniel R. Stinebring,<sup>45</sup> Kevin Stovall,<sup>16</sup> Jerry P. Sun,<sup>30</sup> Joseph K. Swiggum,<sup>14,*</sup> Stephen R. Taylor,<sup>9</sup>
     Jacob E. Turner, 3,4 Michele Vallisneri, 33 Sarah J. Vigeland, 28 Caitlin A. Witt, 3,4 and Kathryn M. Zurek 34
                                                 (The NANOGrav Collaboration)
        <sup>1</sup>X-Ray Astrophysics Laboratory, NASA Goddard Space Flight Center, Code 662, Greenbelt, MD 20771, USA
        <sup>2</sup> Department of Physics and Astronomy, Widener University, One University Place, Chester, PA 19013, USA
     <sup>3</sup> Department of Physics and Astronomy, West Virginia University, P.O. Box 6315, Morgantown, WV 26506, USA
                           <sup>4</sup>Center for Gravitational Waves and Cosmology, West Virginia University,
                                Chestnut Ridge Research Building, Morgantown, WV 26505, USA
                        <sup>5</sup>Department of Physics, Montana State University, Bozeman, MT 59717, USA
                                   <sup>6</sup>Cornell Center for Astrophysics and Planetary Science and
                             Department of Astronomy, Cornell University, Ithaca, NY 14853, USA
                    <sup>7</sup>Cornell Center for Advanced Computing, Cornell University, Ithaca, NY 14853, USA
                               <sup>8</sup>CIFAR Azrieli Global Scholars program, CIFAR, Toronto, Canada
                                 <sup>9</sup>Department of Physics and Astronomy, Vanderbilt University,
                                        2301 Vanderbilt Place, Nashville, TN 37235, USA
                                <sup>10</sup>Station de Radioastronomie de Nancay, Observatoire de Paris,
                               Universite PSL, CNRS, Universite d'Orleans, 18330 Nancau, France
                            <sup>11</sup>FEMTO-ST Institut de recherche, Department of Time and Frequency,
                                       UBFC and CNRS, ENSMM, 25030 Besancon, France
                              <sup>12</sup>Laboratoire de Physique et Chimie de l'Environment et de l'Espace,
                            LPC2E UMR7328, Universite d'Orleans, CNRS, 45071 Orleans, France
     <sup>13</sup>Department of Physics and Astronomy, Franklin & Marshall College, P.O. Box 3003, Lancaster, PA 17604, USA
                              <sup>14</sup>Department of Physics, Lafayette College, Easton, PA 18042, USA
 <sup>15</sup> George Mason University, Fairfax, VA 22030, resident at U.S. Naval Research Laboratory, Washington, D.C. 20375, USA
                   <sup>16</sup>National Radio Astronomy Observatory, 1003 Lopezville Rd., Socorro, NM 87801, USA
                  <sup>17</sup>Department of Physics, Hillsdale College, 33 E. College Street, Hillsdale, MI 49242, USA
                       <sup>18</sup> Eureka Scientific, Inc. 2452 Delmer Street, Suite 100, Oakland, CA 94602-3017
                                      <sup>19</sup>Infinia ML, 202 Rigsbee Avenue, Durham NC, 27701
                         <sup>20</sup>Department of Astronomy, University of Maryland, College Park, MD 20742
        <sup>21</sup>Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771
                               <sup>22</sup>NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
             <sup>23</sup>Department of Physics, McGill University, 3600 University St., Montreal, QC H3A 2T8, Canada
                            <sup>24</sup>Department of Physics and Astronomy, University of British Columbia,
                                    6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada
                    <sup>25</sup> University of Washington Bothell, 18115 Campus Way NE, Bothell, WA 98011, USA
                       <sup>26</sup>Department of Astronomy and National Center for Supercomputing Applications,
                              University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA
                                 <sup>27</sup>McWilliams Center for Cosmology and Department of Physics,
                                     Carnegie Mellon University, Pittsburgh PA, 15213, USA
                                      <sup>28</sup>Center for Gravitation, Cosmology and Astrophysics,
                                   Department of Physics, University of Wisconsin-Milwaukee,
                                            P.O. Box 413, Milwaukee, WI 53201, USA
                                    <sup>29</sup>Center for Interdisciplinary Exploration and Research in
                              Astrophysics (CIERA), Northwestern University, Evanston, IL 60208
                         <sup>30</sup>Department of Physics, Oregon State University, Corvallis, OR 97331, USA
              <sup>31</sup>School of Physics and Astronomy, Rochester Institute of Technology, Rochester, NY 14623, USA
```

<sup>32</sup>Laboratory for Multiwavelength Astrophysics, Rochester Institute of Technology, Rochester, NY 14623, USA

<sup>33</sup> Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA <sup>34</sup> Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, CA <sup>35</sup>Department of Astronomy & Astrophysics, University of Toronto, 50 Saint George Street, Toronto, ON M5S 3H4, Canada <sup>36</sup> Green Bank Observatory, P.O. Box 2, Green Bank, WV 24944, USA <sup>37</sup>Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, New York, New York, 10010, USA <sup>38</sup>Department of Physics, University of Connecticut, 196 Auditorium Road, U-3046, Storrs, CT 06269-3046, USA <sup>39</sup> Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George St., Toronto, ON M5S 3H4, Canada <sup>40</sup>National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA <sup>41</sup>Institute of Physics, Eötvös Loránd University, Pázmány P. s. 1/A, 1117 Budapest, Hungary <sup>42</sup>Space Science Division, Naval Research Laboratory, Washington, DC 20375-5352, USA <sup>43</sup>Department of Astrophysical and Planetary Sciences, University of Colorado, Boulder, CO 80309, USA <sup>44</sup>Centre for Astrophysics and Supercomputing, Swinburne University of Technology, P.O. Box 218, Hawthorn, Victoria 3122, Australia <sup>45</sup>Department of Physics and Astronomy, Oberlin College, Oberlin, OH 44074, USA

We search for a first-order phase transition gravitational wave signal in 45 pulsars from the NANOGrav 12.5 year dataset. We find that the data can be modeled in terms of a strong first order phase transition taking place at temperatures below the electroweak scale. However, we do not observe any strong preference for a phase-transition interpretation of the signal over the standard astrophysical interpretation in terms of supermassive black holes mergers; but we expect to gain additional discriminating power with future datasets, improving the signal to noise ratio and extending the sensitivity window to lower frequencies. An interesting open question is how well gravitational wave observatories could separate such signals.

Introduction — The search for gravitational waves (GWs) spans many orders of magnitude and encapsulates a plethora of source phenomena. At very-low frequencies ( $\sim 1 - 100 \text{ nHz}$ ), pulsar-timing arrays (PTAs; [1-3]) aim to detect GWs through the presence of correlated deviations to radio-pulse arrival times across an ensemble of precisely-timed Milky Way millisecond pulsars. There are three PTA collaborations that currently have decadal-length timing data from an ensemble of pulsars: The North American Nanohertz Observatory for Gravitational Waves (NANOGrav; [4]), the European Pulsar Timing Array (EPTA; [5]), and the Parkes Pulsar Timing Array (PPTA; [6]). These three, in addition to the Indian PTA (InPTA; [7]), are synthesized into the International Pulsar Timing Array (IPTA; Perera et al. 8). There are also emerging efforts in China (CPTA; [9]), as well as some telescope-centered timing programs (MeerKAT; [10]; CHIME; [11]).

The dominant GW signals at such low frequencies frequencies are expected to be from a cosmic population of tightly-bound inspiralling supermassive binary black holes (SMBHBs; [12, 13]), producing an aggregate incoherent signal that we search for as a stochastic GW background (GWB), and also individual binary signals that we attempt to resolve out of this stochastic confusion background. However, other more speculative GW sources in the PTA frequency range include cosmic strings [14, 15], a primordial GWB produced by quan-

tum fluctuations of the gravitational field in the early universe, amplified by inflation [16–18], and cosmological phase transitions [19–23], the latter of which is the subject this study.

The most recent PTA results are from NANOGrav's analysis of 12.5 years of precision timing data from 47 pulsars [24, hereafter NG12], of which 45 exceeded a timing baseline of 3 years and were analysed in a search for a stochastic GWB [25, hereafter NG12gwb]. NANOGrav reported strong evidence for a common-spectrum lowfrequency stochastic process in its array of 45 analyzed pulsars, where  $\sim 10$  of those pulsars are strongly supportive, most are ambivalent, and a few seem to disfavor the process (although not significantly). No evidence for the characteristic inter-pulsar correlation signature imparted by GWs was found. At low frequencies the shape of the characteristic strain spectrum was well matched to a power-law, with an amplitude and slope consistent with theoretical models of SMBHB populations. Under a model that assumes the origin of the GWB is a population of SMBHBs, the median characteristic strain amplitude at a frequency of 1/year is  $1.92 \times 10^{-15}$ . Interpretations of this common-spectrum process as a GWB from SMBHBs have since appeared in the literature, showing that, if it is indeed so, robust evidence of the distinctive inter-pulsar correlations should accrue within the next several years, followed by characterization of the strain spectrum and astrophysical probes of the underlying population [26, 27]. However, the Bayesian posterior probability distributions of the strain-spectrum amplitude and slope are broad enough to entertain a variety of different source interpretations, many of which have since appeared in the literature [e.g. 28–31].

In this Letter we consider gravitational waves produced by first-order cosmological phase transitions, both as an alternative origin of the common process measured in the NANOGrav 12.5 year Dataset [32–39], and as a subdominant signal to that produced by SMBHBs. frequency range to which NANOGrav is sensitive corresponds to phase transitions at temperatures below the electroweak phase transitions of the Standard Model (i.e.  $T \lesssim 100 \,\mathrm{GeV}$ ). This has led many to consider higher frequency GW observatories, such as LISA and LIGO, as the dominant instruments to search for phase transitions. However, phase transitions may occur at much lower temperatures in particular in hidden sectors [40-42]. Hidden sectors/valleys feature rich dynamics, with multiple matter fields and forces, independent of the dynamics of the Standard Model. They appear generically in top-down constructions like string theory, and in some solutions to the so-called hierarchy problem. In many cases, they may be difficult to detect via their particle interactions with the Standard Model, but gravity is an irreducible messenger. In this regard, PTAs provide a powerful complementary probe to the dynamics of hidden sectors already being explored through many terrestrial, astrophysical and cosmological probes (see Ref. [43] for a recent summary).

Previous studies on cosmological first order phase transition in the context of the NANOGrav results were carried out in [33, 37, 44, 45]. Our analysis presents two main novelties compared to these works: first, we properly include the relevant noise sources and discuss the impact of backgrounds (like the one generated by SMB-HBs); second, we discuss how the results are affected by the theoretical uncertainties on the GW spectrum produced by first order phase transitions.

The outline of this Letter is as follows. In the next section we briefly summarize the signature of GWs from the dominant background of SMBH mergers. We then dive into the main subject of this Letter, GWs from a first-order phase transition, where we discuss the relevant parameters characterizing the signal. We then carry out an analysis with the NANOGrav 12.5 year dataset, finding that the data can be modeled in terms of a strong phase transition with a transition temperature around 10 MeV. The dataset and data model for these analyses are exactly as described in NG12 and NG12gwb, respectively. All common processes (whether interpreted as being of SMBHB or phase-transition origin) are modeled within the five lowest sampling frequencies of the array time series, corresponding to  $\sim 2.5-12$  nHz. Finally, we discuss theoretical uncertainties, and compare the PT interpretation of the data to the standard interpretation in terms

	Bubbles [50]	Sound Waves [51]
$\Delta(v_w)$	$\frac{0.48v_w^3}{1 + 5.3v_w^2 + 5v_w^4}$	$0.513v_w$
$\kappa$	$\kappa_\phi$	$\kappa_{ m sw}$
p	2	2
q	2	1
$\mathcal{S}(x)$	$\frac{(a+b)^c}{\left[bx^{-a/c} + ax^{b/c}\right]^c}$	$x^3 \left(\frac{7}{4+3x^2}\right)^{7/2}$
$f_*/\beta$	$\frac{0.35}{1 + 0.07v_w + 0.69v_w^4}$	$\frac{0.536}{v_w}$

TABLE I. Parameters for the gravitational wave spectrum of eq. (4). The values of the parameters (a, b, c) in the spectral shape of the bubble contribution are reported in Table II.

of SMBHB finding no strong preference for one over the other.

GW from SMBHBs mergers — Regardless of origin, the energy density of GWs as a fraction of closure density is related to the GW characteristic strain spectrum by [46]

$$\Omega_{\rm GW}(f) = \frac{2\pi^2}{3H_0^2} f^2 h_c^2(f),\tag{1}$$

where  $H_0$  is the Hubble constant (set here to be 67 km/s/Mpc [47]), and the GWB characteristic strain spectrum  $h_c(f)$  is often described by a power-law function for astrophysical and cosmological sources:

$$h_c(f) = A_{\text{GWB}} \left(\frac{f}{\text{yr}^{-1}}\right)^{\alpha},$$
 (2)

where  $A_{\rm GWB}$  is the amplitude at a reference frequency of 1/year, and  $\alpha$  is an exponent that depends on the origin of the GWB. For a population of inspiraling SMBHBs, this is  $\alpha = -2/3$  [48]. The cross-power spectral density of GW-induced timing deviations between two pulsars a and b can be written as

$$S_{ab}(f) = \Gamma_{ab} \frac{A_{\text{GWB}}^2}{12\pi^2} \left(\frac{f}{\text{yr}^{-1}}\right)^{-\gamma} \text{yr}^3, \qquad (3)$$

where  $\gamma \equiv 3 - 2\alpha = 13/3$  for SMBHBs, and  $\Gamma_{ab}$  is the Hellings-Downs [49] correlation coefficient between pulsar a and pulsar b.

GWs from first-order phase transition — A first-order phase transition (PT) occurs when the true minimum of a potential is separated from a false minimum by a barrier through which a field must locally tunnel. This can occur in either weakly coupled (where a scalar field tunnels) or strongly coupled (where a vacuum condensate corresponds to the scalar field) theory. Such transitions are known to proceed through nucleation of bubbles of

	Envelope	Semi-analytic	Numerical
a	3	1 - 2.2	1.6 - 0.7
b	1	2.6 - 2.9	1.4 - 2.3
c	1.5	1.5 - 3.5	1
$f_*/\beta$	$\frac{0.35}{1 + 0.07v_w + 0.69v_w^2}$	0.1	0.2

TABLE II. Comparison of the bubble spectral shape parameters derived using the envelope and thin wall approximation [50] (left column), the semi-analytic approach of reference [52] (middle column), and lattice simulations [53] (right column). For numerical and semi-analytic results the values of the parameters depend on the choice of the scalar field potential, we report the range of values obtained for the different scalar field potentials considered in the above mentioned works.

true vacuum which, if sufficiently large, expand in the background plasma (still in the false vacuum). Collisions of these bubbles, as well as interactions between the expanding bubble walls and the surrounding plasma, can be efficient sources of GWs.

We characterize the phase transition in terms of four parameters:

- $T_*$  the Universe temperature at which the phase transition takes place.
- $\alpha_*$  the strength of the phase transition, defined as the ratio of the vacuum and relativistic energy density at the time of the phase transition.
- $\beta/H_*$  the bubble nucleation rate in units of the Hubble rate at the time of the phase transition,  $H_*$ .
- $v_w$  the velocity of the bubble walls.

The three main sources of GWs associated with a first-order phase transition are: (i) collisions of bubble walls, (ii) collisions of the sound waves generated in the background plasma by the bubbles expansion, and (iii) turbulence in the plasma generated by expansion and collisions of the sound-wave. However, in this analysis we will not include the turbulence contribution as it usually is subleading compared to the sound-wave one, and also affected by the largest theory uncertanties (see for example [54–56] for recent developments).

The contribution to the total GW spectrum from bubbles and sound waves collisions can be parametrized as [54, 57]

$$h^{2}\Omega(f) = \mathcal{R} \, \Delta(v_{w}) \left(\frac{\kappa \, \alpha_{*}}{1 + \alpha_{*}}\right)^{p} \left(\frac{H_{*}}{\beta}\right)^{q} \mathcal{S}\left(f/f_{*}^{0}\right) , \quad (4)$$

where the prefactor  $\mathcal{R} \simeq 7.69 \times 10^{-5} g_*^{-1/3}$  accounts for the redshift of the GW energy density,  $\mathcal{S}(\cdot)$  parametrizes the spectral shape, and  $\Delta(v_w)$  is a normalization factor which depends on the bubble wall velocity,  $v_w$ . The value

of the peak frequency today,  $f_*^0$ , is related to the value of the peak frequency at emission,  $f_*$ , by:

$$f_*^0 \simeq 1.13 \times 10^{-10} \,\mathrm{Hz} \, \left(\frac{f_*}{\beta}\right) \left(\frac{\beta}{H_*}\right) \left(\frac{T_*}{\mathrm{MeV}}\right) \left(\frac{g_*}{10}\right)^{1/6},$$
(5)

where  $g_*$  denotes the number of relativistic degrees of freedom at the time of the phase transition. The values of the peak frequency at emission, the spectral shape, the normalization factor, and the exponents p and q are reported in Table I for all the production mechanisms considered in this work. Due to the finite lifetime [58, 59] of the sound waves, to derive  $\Omega_{\rm sw}$  eq. (4) needs to be multiplied by a suppression factor  $\Upsilon(\tau_{\rm sw})$  given by [58]:

$$\Upsilon(\tau_{\rm sw}) = 1 - (1 + 2\tau_{\rm sw}H_*)^{-1/2} \tag{6}$$

where the sound-wave lifetime is usually taken to be the timescale for the onset of turbulent behaviors in the plasma [60]:  $\tau_{\rm sw} \approx R_*/\bar{U}_f$ , where the average bubble separation is given by  $R_* = (8\pi)^{1/3}\beta^{-1}{\rm Max}(v_{\rm w},c_s)$  [61], and  $\bar{U}_f^2 \approx 3\kappa_{\rm sw}\alpha/[4(1+\alpha_*)]$  [60].

Generally both the production mechanism contribute to the GW spectrum. However, if the bubble walls interacts with the surrounding plasma most of the energy released in the PT is expected to be transferred to the plasma so that the sound waves (and possibly the turbulence) contribution dominates the GW spectrum. An exception to this scenario is provided by models in which the bubble walls do not interact with the plasma, or by models where the energy released in the PT is large enough that the friction exerted by the plasma is not enough to stop the walls from keep accelerating (runaway scenario). However, determining wether or not the runaway regime is realized is either model dependent or affected by large theoretical uncertainties. Therefore, we perform two separate of analyses. A sound-waveonly (SWO) analysis, where we assume that the runaway regime is not reached and that the sound wave and turbulence contributions dominate the GW spectrum; therefore we set  $\kappa_{\phi} = 0$ , and use the results of reference [62] to derive  $\kappa_{\rm sw}$  as a function of  $v_w$  and  $\alpha_*$ . A bubble-only (BO) analysis, where we assume that the runaway regime is reached and that bubble collisions dominate the GW spectrum; we then set  $v_w = 1$ ,  $\kappa_{sw} = 0$  and  $\kappa_{\phi} = 1$ .

We conclude this section emphasizing that, despite recent progress, large theoretical uncertainties still affect the prediction of the GW signal produced in cosmological phase transitions. To get an idea of the impact that these uncertainties have on our results we will study how the BO analysis is impacted by them. Similar, if not larger, uncertainties affect the sound wave contribution and would impact the results of the SWO analysis.

Assuming that the stress energy density of the expanding bubbles is localized in an infinitesimally thin shell near the bubble wall (thin shell approximation), and that




FIG. 1. In red (blue) the 1- $\sigma$  (68% posterior credible level), and 2- $\sigma$  (95% posterior credible level) contours for the two-dimensional posterior distributions in the  $(T_*, \alpha_*)$  plane obtained in the BO (SWO). The BO analysis has been performed with the spectral shape computed by using the envelope approximation (left panel), semi-analytic results (central panel), and numerical results (right panel). Specifically, we use (a, b, c) = (1, 2.61, 1.5) for the semi-analytic results, and (a, b, c) = (0.7, 2.3, 1) for the numerical results.

it instantaneously decays to zero after two bubbles collide (envelope approximation), the bubble spectral shape can be derived analytically [50, 63]. The spectral shape parameters obtained in this way are reported in the left column of Table II. To go beyond these approximations, 3D lattice simulations are needed. These simulations are extremely expensive given the hierarchy between the large simulation volume needed to include multiple bubbles, and the small lattice spacing needed to resolve the thin walls. Because of the relativistic contraction of the wall width, this separation of scales becomes increasingly large for increasing wall velocities, making it impossible to simulate ultra-relativistic walls. However, the GW spectrum can be simulated at lower velocities and the results extrapolated to larger values. This is the approach taken in Refs. [53, 64], where the authors show that at high frequencies the GW spectrum is much steeper than predicted by the envelope approximation ( $b \sim 1.4 - 2.3$ depending on the form of the scalar field potential). An alternative approach to the problem has been taken by the authors of Refs. [52, 65]. In these works a parametric form for the evolution of the scalar field during bubble collisions is found by using two-bubble simulations. This parametric form is then used in many-bubble simulations to derive the GW spectrum. They also find a steeper high frequency slope ( $b \sim 2.6 - 2.9$ ) compared to the prediction of the envelope approximation. Similar discrepancies are found at low frequencies, where both the numerical and semi-analytic results find a shallower spectrum compared to the envelope approximation (see Tab. II). To probe the theoretical uncertainty associated with the bubble contribution, we will carry out three separate BO analyses utilizing each approach and compare the constraint on the phase transition temperature and

strength.

Results — We now report our results for the BO and SWO analyses. For either of them we consider both the case where the only GW signal is the one produced by the PT, and the one in which the PT signal is superimposed to an astrophysical background produced by SMBHB. This latter analyses will give an indication of how difficult it will be to disentangle a signal from a phase transition from the SMBHB background. The prior distributions for the model parameters of all these analyses, in addition to other noise characterization parameters, are listed in Table III.

The two parameters that we can constrain the most are the transition temperature,  $T_*$ , and the phase transition strength,  $\alpha_*$ . Their 2D posterior distributions for the PT-only searches are shown in Fig. 1. To assess the impact of theoretical uncertainties related to the bubble spectrum, for the BO analysis we report the results obtained by using the three different estimates of the bubble contribution to the GW spectrum described in the previous section (envelope, semi-analytic, and numerical). We can see that at the 1- $\sigma$  (68% posterior credible) level all the searches prefer a strong PT,  $\alpha_* \gtrsim 0.1$ , with low transition temperature,  $T_* \lesssim 10 \, \text{MeV}$ . At 2- $\sigma$  (95% posterior credible) level the posteriors for the semi-analytical and numerical results have support at much higher temperatures, while the envelope results still prefer relatively low values. The preference for small values of  $T_*$  at the 1- $\sigma$  level can be understood by noticing (see Fig. 2) that the data prefer GW spectra that are peaked at frequencies below the NANOGrav sensitivity window (i.e.  $f_*^0 \lesssim 10^{-9} \,\mathrm{Hz}$ ). And, by setting  $\beta/H_* = 1$  in (5), we see that this requirement corresponds to  $T_* \lesssim 10 \,\mathrm{MeV}$ . The low-frequency part of the numerical and semi-analytical

Parameter	Description	Prior	Comments			
White Noise						
$E_k$	EFAC per backend/receiver system	Uniform $[0, 10]$	single-pulsar analysis only			
$Q_k$ [s]	EQUAD per backend/receiver system	log-Uniform $[-8.5, -5]$	single-pulsar analysis only			
$J_k$ [s]	ECORR per backend/receiver system	log-Uniform $[-8.5, -5]$	single-pulsar analysis only			
Red Noise						
$A_{\mathrm{red}}$	red-noise power-law amplitude	log-Uniform $[-20, -11]$	one parameter per pulsar			
$\gamma_{ m red}$	red-noise power-law spectral index	Uniform $[0,7]$	one parameter per pulsar			
Phase Transition						
$T_* [{ m GeV}]$	phase transition temperature	log-Uniform $[-4, 3]$	one parameter for PTA			
$lpha_*$	phase transition strength	log-Uniform $[-1.3, 1]$	one parameter for PTA			
$H_*/\beta$	bubble nucleation rate	log-Uniform $[-2,0]$	one parameter for PTA			
$v_w$	bubble wall velocity	log-Uniform $[-2, 1]$	one parameter for PTA			
Supermassive Black Bole Binaries (SMBHB)						
$A_{\rm GWB}$	common process strain amplitude	log-Uniform $[-18, -14]$	one parameter for PTA			
$\gamma_{ m GWB}$	common process power-law spectral index	delta function ( $\gamma_{\text{GWB}} = 13/3$ )	fixed			

TABLE III. Priors distributions for the parameters used in all the analyses in this work. The prior for the bubble wall velocity reported in this table is the one used for the SWO analysis, for the BO analyses we use  $v_w = 1$  as explained in the text.

GW spectra is shallow enough that, at the  $2-\sigma$  level, the data can be fitted also by spectra with peak frequencies above the NANOGrav sensitivity window. The same is not true for the envelope results, which have a much steeper low-frequency spectrum; this is the reason why the  $2-\sigma$  levels of the envelope results deviate substantially from the other two.

In Fig. 2 we show the GWB spectrum predicted for the maximum likelihood parameters of PT-only searches. To better illustrate our results, and how the different parameters and theoretical uncertainties affect the GWB spectrum, we release an interactive version of Fig. 2 at this link.

To understand how the inclusion of the SMBHB background affects our results, in Fig. 3 we show the posterior for the parameters  $\alpha_*$  and  $A_{\rm GWB}$  obtained in the PT+SMBHB search. As expected, with the inclusion of the SMBHB background, the posteriors for  $\alpha_*$  stretch to lower values where most of the signal is provided by the SMBHB contribution. The Bayesian Information Criterion (BIC) [66], defined to be BIC =  $k \ln n - 2 \ln \hat{\mathcal{L}}$  where n=5 is the number of data points in the frequency space, k is the number of parameters in the model and  $\hat{\mathcal{L}}$  is the maximum likelihood, is also computed. For the BO searches, the differences in BIC between the PT+SMBHB and SMBHB only searches are found to be -0.92, 3.04 and 1.89 for the envelope, semi-analytic

and numerical results respectively; similarly the BIC differences between the PT-only and SMBHB-only searches are -1.82, -3.18, -1.28. For the SWO analysis, the difference in BIC between the PT+SMBHB and SMBHB only searches is -4.56, while we get -2.19 for the difference between the PT-only and SMBHB-only searches. We can then conclude that that the PT+SMBHB and PT-only models were neither strongly favored nor disfavored compared to the SMBHB only model [67].

A complete set of posteriors for the parameters of the PT-only searches (derived by using the semi-analytic spectrum for the BO analysis) are shown in Fig. 4. As noted previously, at 1- $\sigma$  level the data prefer a strongly first-order phase transition ( $\alpha_* \gtrsim 0.1$ ) taking place at temperature  $T_* \lesssim 10 \,\mathrm{MeV}$ ; while no strong constraints on  $v_w$  or  $H_*/\beta$  is observed. We can also notice that the higher values of  $T_*$  allowed in the 2- $\sigma$  region are accompanied by slower nucleation rates (large  $H_*/\beta$ ). We should caution, however, that numerical simulation have been performed for phase transition strengths up to  $\alpha_* \sim 0.5$ [68], and that our results for  $\alpha_* \gtrsim 0.5$  are derived by extrapolating the results of these simulations. A similar remark should be made for  $H_*/\beta$ , numerical simulations with values of this parameter close to unity have not been performed yet.

Given the low value of  $T_*$ , and the strong constraints on new physics at such low scales, we expect the phase transition to take place in a dark sector with only feeble interactions with the Standard Model (SM). In order to be consistent with the Hubble parameter constraints during the era of Big Bang Nucleosynthesis (BBN) [69], the energy of this dark sector must be transferred to the

<sup>&</sup>lt;sup>1</sup> The posterior stops around  $\alpha_* \simeq 0.05$  because of choice of the prior, otherwise it would extend down to  $\alpha_* \sim 0$ .




FIG. 2. Maximum likelihood GWB fractional energy-density spectrum for the BO (red) and SWO (blue) analyses compared with the marginalized posterior for the free power spectrum (independent per-frequency characterization; red violin plot) derived in NG12gwb. For the BO analysis we show the results derived by using the envelope (solid line), semi-analytic (dashed), and numerical (dot-dashed) spectral shapes. For the BO analyses the values of  $(\alpha_*, T_*)$  for these maximum likelihood spectra are  $(0.28, 0.7\,\mathrm{MeV})$  for the envelope results,  $(1.2, 3.4\,\mathrm{MeV})$  for the semi-analytic results, and  $(0.13, 14.1\,\mathrm{MeV})$  for the numerical results. While for the SO analysis we get  $(6.0, 0.32\,\mathrm{MeV})$ .

SM before the onset of BBN at  $T \sim 1 \,\mathrm{MeV}$ . This leaves an allowed range of values for the transition temperature given by  $T_* \sim 1 \,\mathrm{MeV} - 100 \,\mathrm{GeV}$ . The next data release, which adds multiple years of observations and extends the the sensitivity window to lower frequency, should begin to resolve the peak of the spectrum or additionally shrink the range of allowed values for  $T_*$ .

Conclusions — We performed a search for a stochastic gravitational wave background from first-order phase transitions in the 12.5 year NANOGrav dataset. While previous NANOGrav analysis found no evidence yet for the inter-pulsar correlation signature of a GWB, the evidence for a common-spectrum process was significant. We found that the data can be modeled by a strong  $(\alpha_* > 0.1)$  phase transition taking place at temperatures below the electroweak scale. However, the data do not show any strong preference between an SMBHB and a PT generated signal, but we expect to gain additional discriminating power with future datasets, improving the signal to noise ratio and extending the sensitivity window to lower frequencies. In particular, data from the International Pulsar Timing Array will allow the baseline of observations to be significantly extended, and the num-




FIG. 3. 1- $\sigma$  (68% posterior credible level), and 2- $\sigma$  (95% posterior credible level) contours for the parameters  $A_{\rm GWB}$  and  $\alpha_*$  in the PT+SMBHB search. In red (blue) the results for the BO (SWO) analyses. In this figure we have used the semi-analytic results for the bubble spectrum. The posteriors do not extend to lower values of  $\alpha_*$  because of our choice for the  $\alpha_*$  prior: log-Uniform [-1.3, 1].

ber of monitored pulsars to be greatly expanded. The present quality of the data is such that our results are not strongly affected by theoretical uncertainties on the GW spectral shape. However, methodological improvements on determining the origin of the GWB spectrum will be needed for future datasets in order to separate the signal from a first-order PT from the SMBHB background, as well as to constrain the microscopic origins of the PT.

Author contributions — An alphabetical-order author list was used for this paper in recognition of the fact that a large, decade timescale project such as NANOGrav is necessarily the result of the work of many people. All authors contributed to the activities of the NANOGrav collaboration leading to the work presented here, and reviewed the manuscript, text, and figures prior to the paper's submission. Additional specific contributions to this paper are as follows. ZA, HB, PRB, HTC, MED, PBD, TD, JAE, RDF, ECF, EF, NG-D, PAG, DCG, MLJ, MTL, DRL, RSL, JL, MAM, CN, DJN, TTP, NSP, SMR, KS, IHS, RS, JKS, RS and SJV developed the 12.5-year dataset through a combination of observations, arrival time calculations, data checks and refinements, and timing model development and analysis; additional specific contributions to the dataset are summarized in NG12. KZ and SRT coordinated the writing of the paper. VL and AM performed all analyses presented in




FIG. 4. Corner plot showing the 1D and 2D posterior distributions for the parameters of the PT-only search. In red (blue) the results for the BO (SWO) analyses. In deriving these results we have used the semi-analytic bubble spectral shape with (a, b, c) = (1, 2.61, 1.5).

this paper. KZ, SRT, AM, and VL wrote the paper and collected the bibliography.

Acknowledgments — This work has been carried out by the NANOGrav collaboration, which is part of the International Pulsar Timing Array. The NANOGrav project receives support from National Science Foundation (NSF) Physics Frontiers Center award number 1430284. The Arecibo Observatory is a facility of the

NSF operated under cooperative agreement (#AST-1744119) by the University of Central Florida (UCF) in alliance with Universidad Ana G. Méndez (UAGM) and Yang Enterprises (YEI), Inc. The Green Bank Observatory is a facility of the NSF operated under cooperative agreement by Associated Universities, Inc. The National Radio Astronomy Observatory is a facility of the NSF operated under cooperative agreement by As-

sociated Universities, Inc. A majority of the computational work was performed on the Caltech High Performance Cluster, partially supported by a grant from the Gordon and Betty Moore Foundation.. This work made use of the Super Computing System (Spruce Knob) at WVU, which are funded in part by the National Science Foundation EPSCoR Research Infrastructure Improvement Cooperative Agreement #1003907, the state of West Virginia (WVEPSCoR via the Higher Education Policy Commission) and WVU. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. Portions of this work performed at NRL were supported by Office of Naval Research 6.1 funding. The Flatiron Institute is supported by the Simons Foundation. Pulsar research at UBC is supported by an NSERC Discovery Grant and by the Canadian Institute for Advanced Research. SRT acknowledges support from NSF grant AST-#2007993, and a Dean's Faculty Fellowship from Vanderbilt University's College of Arts & Science. JS and MV acknowledge support from the JPL RTD program. SBS acknowledges support for this work from NSF grants #1458952 and #1815664. SBS is a CIFAR Azrieli Global Scholar in the Gravity and the Extreme Universe program. TTP acknowledges support from the MTA-ELTE Extragalactic Astrophysics Research Group, funded by the Hungarian Academy of Sciences (Magyar Tudományos Akadémia), that was used during the development of this research. TD and ML acknowledge NSF AAG award number 2009468. This work is supported in part by NASA under award number 80GSFC17M0002. VL, AM and KZ are supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award Number DE-SC0021431 and a Simons Investigator award

Facilities — Arecibo, GBT
Software — ENTERPRISE [70],
enterprise\_extensions [71], HASASIA [72], libstempo
[73], matplotlib [74], PTMCMC [75], tempo [76], tempo2
[77], PINT [78]

- \* NANOGrav Physics Frontiers Center Postdoctoral Fellow
- R. S. Foster and D. C. Backer, Constructing a Pulsar Timing Array, ApJ 361, 300 (1990).
- [2] S. Detweiler, Pulsar timing measurements and the search for gravitational waves, ApJ 234, 1100 (1979).
- [3] M. V. Sazhin, Opportunities for detecting ultralong gravitational waves, Soviet Ast. 22, 36 (1978).
- [4] S. Ransom, A. Brazier, S. Chatterjee, T. Cohen, J. M. Cordes, M. E. DeCesar, P. B. Demorest, J. S. Hazboun, M. T. Lam, R. S. Lynch, M. A. McLaughlin, S. M. Ransom, X. Siemens, S. R. Taylor, and S. J. Vigeland, The NANOGrav Program for Gravitational Waves and

- Fundamental Physics, in BAAS, Vol. 51 (2019) p. 195, arXiv:1908.05356 [astro-ph.IM].
- [5] G. Desvignes, R. N. Caballero, L. Lentati, J. P. W. Verbiest, D. J. Champion, B. W. Stappers, G. H. Janssen, P. Lazarus, S. Osłowski, S. Babak, C. G. Bassa, P. Brem, M. Burgay, I. Cognard, J. R. Gair, E. Graikou, L. Guillemot, J. W. T. Hessels, A. Jessner, C. Jordan, R. Karuppusamy, M. Kramer, A. Lassus, K. Lazaridis, K. J. Lee, K. Liu, A. G. Lyne, J. McKee, C. M. F. Mingarelli, D. Perrodin, A. Petiteau, A. Possenti, M. B. Purver, P. A. Rosado, S. Sanidas, A. Sesana, G. Shaifullah, R. Smits, S. R. Taylor, G. Theureau, C. Tiburzi, R. van Haasteren, and A. Vecchio, High-precision timing of 42 millisecond pulsars with the European Pulsar Timing Array, MNRAS 458, 3341 (2016), arXiv:1602.08511 [astro-ph.HE].
- [6] M. Kerr, D. J. Reardon, G. Hobbs, R. M. Shannon, R. N. Manchester, S. Dai, C. J. Russell, S. B. Zhang, W. van Straten, S. Osłowski, A. Parthasarathy, R. Spiewak, M. Bailes, N. D. R. Bhat, A. D. Cameron, W. A. Coles, J. Dempsey, X. Deng, B. Goncharov, J. F. Kaczmarek, M. J. Keith, P. D. Lasky, M. E. Lower, B. Preisig, J. M. Sarkissian, L. Toomey, H. Wang, J. Wang, L. Zhang, and X. Zhu, The Parkes Pulsar Timing Array Project: Second data release, arXiv e-prints, arXiv:2003.09780 (2020), arXiv:2003.09780 [astro-ph.IM].
- [7] B. C. Joshi, P. Arumugasamy, M. Bagchi, D. Bandyopadhyay, A. Basu, N. Dhand a Batra, S. Bethapudi, A. Choudhary, K. De, L. Dey, A. Gopakumar, Y. Gupta, M. A. Krishnakumar, Y. Maan, P. K. Manoharan, A. Naidu, R. Nandi, D. Pathak, M. Surnis, and A. Susobhanan, Precision pulsar timing with the ORT and the GMRT and its applications in pulsar astrophysics, Journal of Astrophysics and Astronomy 39, 51 (2018).
- [8] B. B. P. Perera, M. E. DeCesar, P. B. Demorest, M. Kerr. L. Lentati, D. J. Nice, S. Osłowski, S. M. Ransom, M. J. Keith, Z. Arzoumanian, M. Bailes, P. T. Baker, C. G. Bassa, N. D. R. Bhat, A. Brazier, M. Burgay, S. Burke-Spolaor, R. N. Caballero, D. J. Champion, S. Chatterjee, S. Chen, I. Cognard, J. M. Cordes, K. Crowter, S. Dai, G. Desvignes, T. Dolch, R. D. Ferdman, E. C. Ferrara, E. Fonseca, J. M. Goldstein, E. Graikou, L. Guillemot, J. S. Hazboun, G. Hobbs, H. Hu, K. Islo, G. H. Janssen, R. Karuppusamy, M. Kramer, M. T. Lam, K. J. Lee, K. Liu, J. Luo, A. G. Lyne, R. N. Manchester, J. W. McKee, M. A. McLaughlin, C. M. F. Mingarelli, A. P. Parthasarathy, T. T. Pennucci, D. Perrodin, A. Possenti, D. J. Reardon, C. J. Russell, S. A. Sanidas, A. Sesana, G. Shaifullah, R. M. Shannon, X. Siemens, J. Simon, R. Spiewak, I. H. Stairs, B. W. Stappers, J. K. Swiggum, S. R. Taylor, G. Theureau, C. Tiburzi, M. Vallisneri, A. Vecchio, J. B. Wang, S. B. Zhang, L. Zhang, W. W. Zhu, and X. J. Zhu, The International Pulsar Timing Array: second data release, MNRAS 490, 4666 (2019), arXiv:1909.04534 [astro-ph.HE].
- [9] K. J. Lee, Prospects of Gravitational Wave Detection Using Pulsar Timing Array for Chinese Future Telescopes, in Frontiers in Radio Astronomy and FAST Early Sciences Symposium 2015, Astronomical Society of the Pacific Conference Series, Vol. 502, edited by L. Qain and D. Li (2016) p. 19.
- [10] M. Bailes, E. Barr, N. D. R. Bhat, J. Brink, S. Buchner, M. Burgay, F. Camilo, D. J. Champion, J. Hessels, G. H. Janssen, A. Jameson, S. Johnston, A. Karastergiou, R. Karuppusamy, V. Kaspi, M. J. Keith, M. Kramer,

- M. A. McLaughlin, K. Moodley, S. Oslowski, A. Possenti, S. M. Ransom, F. A. Rasio, J. Sievers, M. Serylak, B. W. Stappers, I. H. Stairs, G. Theureau, W. van Straten, P. Weltevrede, and N. Wex, MeerTime the MeerKAT Key Science Program on Pulsar Timing, arXiv e-prints , arXiv:1803.07424 (2018), arXiv:1803.07424 [astro-ph.IM].
- [11] C. Ng, Pulsar science with the CHIME telescope, in Pulsar Astrophysics the Next Fifty Years, IAU Symposium, Vol. 337, edited by P. Weltevrede, B. B. P. Perera, L. L. Preston, and S. Sanidas (2018) pp. 179–182, arXiv:1711.02104 [astro-ph.IM].
- [12] A. Sesana, F. Haardt, P. Madau, and M. Volonteri, Low-Frequency Gravitational Radiation from Coalescing Massive Black Hole Binaries in Hierarchical Cosmologies, ApJ 611, 623 (2004), astro-ph/0401543.
- [13] S. Burke-Spolaor, S. R. Taylor, M. Charisi, T. Dolch, J. S. Hazboun, A. M. Holgado, L. Z. Kelley, T. J. W. Lazio, D. R. Madison, N. McMann, C. M. F. Mingarelli, A. Rasskazov, X. Siemens, J. J. Simon, and T. L. Smith, The astrophysics of nanohertz gravitational waves, A&A Rev. 27, 5 (2019), arXiv:1811.08826 [astroph.HE].
- [14] X. Siemens, V. Mandic, and J. Creighton, Gravitational-Wave Stochastic Background from Cosmic Strings, Phys. Rev. Lett. 98, 111101 (2007), arXiv:astro-ph/0610920 [astro-ph].
- [15] J. J. Blanco-Pillado, K. D. Olum, and X. Siemens, New limits on cosmic strings from gravitational wave observation, Physics Letters B 778, 392 (2018), arXiv:1709.02434 [astro-ph.CO].
- [16] L. P. Grishchuk, Amplification of gravitational waves in an isotropic universe, Soviet Journal of Experimental and Theoretical Physics 40, 409 (1975).
- [17] P. D. Lasky, C. M. F. Mingarelli, T. L. Smith, J. T. Giblin, E. Thrane, D. J. Reardon, R. Caldwell, M. Bailes, N. D. R. Bhat, S. Burke-Spolaor, S. Dai, J. Dempsey, G. Hobbs, M. Kerr, Y. Levin, R. N. Manchester, S. Osłowski, V. Ravi, P. A. Rosado, R. M. Shannon, R. Spiewak, W. van Straten, L. Toomey, J. Wang, L. Wen, X. You, and X. Zhu, Gravitational-Wave Cosmology across 29 Decades in Frequency, Physical Review X 6, 011035 (2016), arXiv:1511.05994 [astro-ph.CO].
- [18] S. Vagnozzi, Implications of the NANOGrav results for inflation, Mon. Not. Roy. Astron. Soc. 502, L11 (2021), arXiv:2009.13432 [astro-ph.CO].
- [19] J. Winicour, Gravitational Radiation from Relativistic Phase Transitions, ApJ 182, 919 (1973).
- [20] C. J. Hogan, Gravitational radiation from cosmological phase transitions, Mon. Not. Roy. Astron. Soc. 218, 629 (1986).
- [21] D. V. Deryagin, D. Y. Grigoriev, V. A. Rubakov, and M. V. Sazhin, Possible Anisotropic Phases in the Early Universe and Gravitational Wave Background, Mod. Phys. Lett. A 1, 593 (1986).
- [22] C. Caprini, R. Durrer, and X. Siemens, Detection of gravitational waves from the QCD phase transition with pulsar timing arrays, Phys. Rev. D 82, 063511 (2010), arXiv:1007.1218 [astro-ph.CO].
- [23] A. Kobakhidze, C. Lagger, A. Manning, and J. Yue, Gravitational waves from a supercooled electroweak phase transition and their detection with pulsar timing arrays, The European Physical Journal C 77 (2017).
- [24] M. F. Alam, Z. Arzoumanian, P. T. Baker, H. Blumer,

- K. E. Bohler, A. Brazier, P. R. Brook, S. Burke-Spolaor, K. Caballero, R. S. Camuccio, R. L. Chamberlain, S. Chatterjee, J. M. Cordes, N. J. Cornish, F. Crawford, H. T. Cromartie, M. E. DeCesar, P. B. Demorest, T. Dolch, J. A. Ellis, R. D. Ferdman, E. C. Ferrara, W. Fiore, E. Fonseca, Y. Garcia, N. Garver-Daniels, P. A. Gentile, D. C. Good, J. A. Gusdorff, D. Halmrast, J. Hazboun, K. Islo, R. J. Jennings, C. Jessup, M. L. Jones, A. R. Kaiser, D. L. Kaplan, L. Z. Kelley, J. Shapiro Key, M. T. Lam, T. J. W. Lazio, D. R. Lorimer, J. Luo, R. S. Lynch, D. Madison, K. Maraccini, M. A. McLaughlin, C. M. F. Mingarelli, C. Ng, B. M. X. Nguyen, D. J. Nice, T. T. Pennucci, N. S. Pol, J. Ramette, S. M. Ransom, P. S. Ray, B. J. Shapiro-Albert, X. Siemens, J. Simon, R. Spiewak, I. H. Stairs, D. R. Stinebring, K. Stovall, J. K. Swiggum, S. R. Taylor, M. Tripepi, M. Vallisneri, S. J. Vigeland, C. A. Witt, and W. Zhu, The NANOGrav 12.5-year Data Set: Observations and Narrowband Timing of 47 Millisecond Pulsars, arXiv e-prints, arXiv:2005.06490 (2020), arXiv:2005.06490 [astro-ph.HE].
- [25] Z. Arzoumanian, P. T. Baker, H. Blumer, B. Bécsy, A. Brazier, P. R. Brook, S. Burke-Spolaor, S. Chatterjee, S. Chen, J. M. Cordes, N. J. Cornish, F. Crawford, H. T. Cromartie, M. E. Decesar, P. B. Demorest, T. Dolch, J. A. Ellis, E. C. Ferrara, W. Fiore, E. Fonseca, N. Garver-Daniels, P. A. Gentile, D. C. Good, J. S. Hazboun, A. M. Holgado, K. Islo, R. J. Jennings, M. L. Jones, A. R. Kaiser, D. L. Kaplan, L. Z. Kelley, J. S. Key, N. Laal, M. T. Lam, T. J. W. Lazio, D. R. Lorimer, J. Luo, R. S. Lynch, D. R. Madison, M. A. McLaughlin, C. M. F. Mingarelli, C. Ng, D. J. Nice, T. T. Pennucci, N. S. Pol, S. M. Ransom, P. S. Ray, B. J. Shapiro-Albert, X. Siemens, J. Simon, R. Spiewak, I. H. Stairs, D. R. Stinebring, K. Stovall, J. P. Sun. J. K. Swiggum, S. R. Taylor, J. E. Turner, M. Vallisneri, S. J. Vigeland, C. A. Witt, and Nanograv Collaboration, The NANOGrav 12.5 yr Data Set: Search for an Isotropic Stochastic Gravitational-wave Background, ApJ 905, L34 (2020), arXiv:2009.04496 [astro-ph.HE].
- [26] H. Middleton, A. Sesana, S. Chen, A. Vecchio, W. Del Pozzo, and P. A. Rosado, Massive black hole binary systems and the NANOGrav 12.5 yr results, MNRAS 502, L99 (2021), arXiv:2011.01246 [astro-ph.HE].
- [27] N. S. Pol, S. R. Taylor, L. Z. Kelley, S. J. Vigeland, J. Simon, S. Chen, Z. Arzoumanian, P. T. Baker, B. Bécsy, A. Brazier, P. R. Brook, S. Burke-Spolaor, S. Chatterjee, J. M. Cordes, N. J. Cornish, F. Crawford, H. T. Cromartie, M. E. DeCesar, P. B. Demorest, T. Dolch, E. C. Ferrara, W. Fiore, E. Fonseca, N. Garver-Daniels, D. C. Good, J. S. Hazboun, R. J. Jennings, M. L. Jones, A. R. Kaiser, D. L. Kaplan, J. Shapiro Key, M. T. Lam, T. J. W. Lazio, J. Luo, R. S. Lynch, D. R. Madison, A. McEwen, M. A. McLaughlin, C. M. F. Mingarelli, C. Ng, D. J. Nice, T. T. Pennucci, S. M. Ransom, P. S. Ray, B. J. Shapiro-Albert, X. Siemens, I. H. Stairs, D. R. Stinebring, J. K. Swiggum, M. Vallisneri, H. Wahl, and C. A. Witt, Astrophysics Milestones For Pulsar Timing Array Gravitational Wave Detection, arXiv e-prints, arXiv:2010.11950 (2020), arXiv:2010.11950 [astro-ph.HE].
- [28] V. Vaskonen and H. Veermäe, Did nanograv see a signal from primordial black hole formation?, Phys. Rev. Lett. 126, 051303 (2021).

- [29] V. De Luca, G. Franciolini, and A. Riotto, Nanograv data hints at primordial black holes as dark matter, Phys. Rev. Lett. 126, 041303 (2021).
- [30] J. Ellis and M. Lewicki, Cosmic string interpretation of nanograv pulsar timing data, Phys. Rev. Lett. 126, 041304 (2021).
- [31] S. Blasi, V. Brdar, and K. Schmitz, Has nanograv found first evidence for cosmic strings?, Phys. Rev. Lett. 126, 041305 (2021).
- [32] A. Brandenburg, E. Clarke, Y. He, and T. Kahniashvili, Can we observe the QCD phase transition-generated gravitational waves through pulsar timing arrays?, arXiv e-prints, arXiv:2102.12428 (2021), arXiv:2102.12428 [astro-ph.CO].
- [33] A. Neronov, A. R. Pol, C. Caprini, and D. Semikoz, NANOGrav signal from magnetohydrodynamic turbulence at the QCD phase transition in the early Universe, Phys. Rev. D 103, L041302 (2021), arXiv:2009.14174 [astro-ph.CO].
- [34] S.-L. Li, L. Shao, P. Wu, and H. Yu, NANOGrav Signal from First-Order Confinement/Deconfinement Phase Transition in Different QCD Matters, arXiv e-prints, arXiv:2101.08012 (2021), arXiv:2101.08012 [astro-ph.CO].
- [35] B. Barman, A. Dutta Banik, and A. Paul, Implications of NANOGrav results and UV freeze-in in a fastexpanding Universe, arXiv e-prints, arXiv:2012.11969 (2020), arXiv:2012.11969 [astro-ph.CO].
- [36] K. T. Abe, Y. Tada, and I. Ueda, Induced gravitational waves as a cosmological probe of the sound speed during the QCD phase transition, arXiv e-prints, arXiv:2010.06193 (2020), arXiv:2010.06193 [astro-ph.CO].
- [37] W. Ratzinger and P. Schwaller, Whispers from the dark side: Confronting light new physics with NANOGrav data, arXiv e-prints, arXiv:2009.11875 (2020), arXiv:2009.11875 [astro-ph.CO].
- [38] A. Addazi, Y.-F. Cai, Q. Gan, A. Marciano, and K. Zeng, NANOGrav results and Dark First Order Phase Transitions, arXiv e-prints, arXiv:2009.10327 (2020), arXiv:2009.10327 [hep-ph].
- [39] Y. Nakai, M. Suzuki, F. Takahashi, and M. Yamada, Gravitational Waves and Dark Radiation from Dark Phase Transition: Connecting NANOGrav Pulsar Timing Data and Hubble Tension, arXiv e-prints, arXiv:2009.09754 (2020), arXiv:2009.09754 [astro-ph.CO].
- [40] M. J. Strassler and K. M. Zurek, Echoes of a hidden valley at hadron colliders, Phys. Lett. B 651, 374 (2007), arXiv:hep-ph/0604261.
- [41] Z. Chacko, L. J. Hall, and Y. Nomura, Acceleressence: dark energy from a phase transition at the seesaw scale, JCAP 10, 011, arXiv:astro-ph/0405596.
- [42] P. Schwaller, Gravitational Waves from a Dark Phase Transition, Phys. Rev. Lett. 115, 181101 (2015), arXiv:1504.07263 [hep-ph].
- [43] M. Battaglieri, A. Belloni, A. Chou, P. Cushman, B. Echenard, R. Essig, J. Estrada, J. L. Feng, B. Flaugher, P. J. Fox, P. Graham, C. Hall, R. Harnik, J. Hewett, J. Incandela, E. Izaguirre, D. McKinsey, M. Pyle, N. Roe, G. Rybka, P. Sikivie, T. M. P. Tait, N. Toro, R. Van De Water, N. Weiner, K. Zurek, E. Adelberger, A. Afanasev, D. Alexander, J. Alexander, V. Cristian Antochi, D. M. Asner, H. Baer, D. Baner-
- jee, E. Baracchini, P. Barbeau, J. Barrow, N. Bastidon, J. Battat, S. Benson, A. Berlin, M. Bird, N. Blinov, K. K. Boddy, M. Bondi, W. M. Bonivento, M. Boulay, J. Boyce, M. Brodeur, L. Broussard, R. Budnik, P. Bunting, M. Caffee, S. S. Caiazza, S. Campbell, T. Cao, G. Carosi, M. Carpinelli, G. Cavoto, A. Celentano, J. Hyeok Chang, S. Chattopadhyay, A. Chavarria, C.-Y. Chen, K. Clark, J. Clarke, O. Colegrove, J. Coleman, D. Cooke, R. Cooper, M. Crisler, P. Crivelli, F. D'Eramo, D. D'Urso, E. Dahl, W. Dawson, M. De Napoli, R. De Vita, P. DeNiverville, S. Derenzo, A. Di Crescenzo, E. Di Marco, K. R. Dienes, M. Diwan, D. Handiipondola Dongwi, A. Drlica-Wagner, S. Ellis, A. Chigbo Ezeribe, G. Farrar, F. Ferrer, E. Figueroa-Feliciano, A. Filippi, G. Fiorillo, B. Fornal, A. Freyberger, C. Frugiuele, C. Galbiati, I. Galon, S. Gardner, A. Geraci, G. Gerbier, M. Graham, E. Gschwendtner, C. Hearty, J. Heise, R. Henning, R. J. Hill, D. Hitlin, Y. Hochberg, J. Hogan, M. Holtrop, Z. Hong, T. Hossbach, T. B. Humensky, P. Ilten, K. Irwin, J. Jaros, R. Johnson, M. Jones, Y. Kahn, N. Kalantarians, M. Kaplinghat, R. Khatiwada, S. Knapen, M. Kohl, C. Kouvaris, J. Kozaczuk, G. Krnjaic, V. Kubarovsky, E. Kuflik, A. Kusenko, R. Lang, K. Leach, T. Lin, M. Lisanti, J. Liu, K. Liu, M. Liu, D. Loomba, J. Lykken, K. Mack, J. Mans, H. Maris, T. Markiewicz, L. Marsicano, C. J. Martoff, G. Mazzitelli, C. McCabe, S. D. Mc-Dermott, A. McDonald, B. McKinnon, D. Mei, T. Melia, G. A. Miller, K. Miuchi, S. M. P. Nazeer, O. Moreno, V. Morozov, F. Mouton, H. Mueller, A. Murphy, R. Neilson, T. Nelson, C. Neu, Y. Nosochkov, C. O'Hare, N. Oblath, J. Orrell, J. Ouellet, S. Pastore, S. Paul, M. Perelstein, A. Peter, N. Phan, N. Phinney, M. Pivovaroff, A. Pocar, M. Pospelov, J. Pradler, P. Privitera, S. Profumo, M. Raggi, S. Rajendran, N. Randazzo, T. Raubenheimer, C. Regenfus, A. Renshaw, A. Ritz, T. Rizzo, L. Rosenberg, A. Rubbia, B. Rybolt, T. Saab, B. R. Safdi, E. Santopinto, A. Scarff, M. Schneider, P. Schuster, G. Seidel, H. Sekiya, I. Seong, G. Simi, V. Sipala, T. Slatyer, O. Slone, P. F. Smith, J. Smolinsky, D. Snowden-Ifft, M. Solt, A. Sonnenschein, P. Sorensen, N. Spooner, B. Srivastava, I. Stancu, L. Strigari, J. Strube, A. O. Sushkov, M. Szydagis, P. Tanedo, D. Tanner, R. Tayloe, W. Terrano, J. Thaler, B. Thomas, B. Thorpe, T. Thorpe, J. Tiffenberg, N. Tran, M. Trovato, C. Tully, T. Tyson, T. Vachaspati, S. Vahsen, K. van Bibber, J. Vandenbroucke, A. Villano, T. Volansky, G. Wang, T. Ward, W. Wester, A. Whitbeck, D. A. Williams, M. Wing, L. Winslow, B. Wojtsekhowski, H.-B. Yu, S.-S. Yu, T.-T. Yu, X. Zhang, Y. Zhao, and Y.-M. Zhong, US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report, arXiv eprints, arXiv:1707.04591 (2017), arXiv:1707.04591 [hepph].
- [44] S.-L. Li, L. Shao, P. Wu, and H. Yu, NANOGrav Signal from First-Order Confinement/Deconfinement Phase Transition in Different QCD Matters, (2021), arXiv:2101.08012 [astro-ph.CO].
- [45] L. Bian, R.-G. Cai, J. Liu, X.-Y. Yang, and R. Zhou, On the gravitational wave sources from the NANOGrav 12.5-yr data, (2020), arXiv:2009.13893 [astro-ph.CO].
- [46] C. J. Moore, R. H. Cole, and C. P. L. Berry, Gravitational-wave sensitivity curves, Classical and Quantum Gravity 32, 015014 (2015), arXiv:1408.0740

[gr-qc].

- [47] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J. P. Bernard, M. Bersanelli, P. Bielewicz, J. J. Bock, J. R. Bond, J. Borrill, F. R. Bouchet, F. Boulanger, M. Bucher, C. Burigana, R. C. Butler, E. Calabrese, J. F. Cardoso, J. Carron, A. Challinor, H. C. Chiang, J. Chluba, L. P. L. Colombo, C. Combet, D. Contreras, B. P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille, J. M. Delouis, E. Di Valentino, J. M. Diego, O. Doré, M. Douspis, A. Ducout, X. Dupac, S. Dusini, G. Efstathiou, F. Elsner, T. A. Enßlin, H. K. Eriksen, Y. Fantaye, M. Farhang, J. Fergusson, R. Fernandez-Cobos, F. Finelli, F. Forastieri, M. Frailis, A. A. Fraisse, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R. T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K. M. Górski, S. Gratton, A. Gruppuso, J. E. Gudmundsson, J. Hamann, W. Handley, F. K. Hansen, D. Herranz, S. R. Hildebrandt, E. Hivon, Z. Huang, A. H. Jaffe, W. C. Jones, A. Karakci, E. Keihänen, R. Keskitalo, K. Kiiveri, J. Kim, T. S. Kisner, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J. M. Lamarre, A. Lasenby, M. Lattanzi, C. R. Lawrence, M. Le Jeune, P. Lemos, J. Lesgourgues, F. Levrier, A. Lewis, M. Liguori, P. B. Lilje, M. Lilley, V. Lindholm, M. López-Caniego, P. M. Lubin, Y. Z. Ma, J. F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P. G. Martin, M. Martinelli, E. Martínez-González, S. Matarrese, N. Mauri, J. D. McEwen, P. R. Meinhold, A. Melchiorri, A. Mennella, M. Migliaccio, M. Millea, S. Mitra, M. A. Miville-Deschênes, D. Molinari, L. Montier, G. Morgante, A. Moss, P. Natoli, H. U. Nørgaard-Nielsen, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, H. V. Peiris, F. Perrotta, V. Pettorino, F. Piacentini, L. Polastri, G. Polenta, J. L. Puget, J. P. Rachen, M. Reinecke, M. Remazeilles, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J. A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, E. P. S. Shellard, C. Sirignano, G. Sirri, L. D. Spencer, R. Sunyaev, A. S. Suur-Uski, J. A. Tauber, D. Tavagnacco, M. Tenti, L. Toffolatti, M. Tomasi, T. Trombetti, L. Valenziano, J. Valiviita, B. Van Tent, L. Vibert, P. Vielva, F. Villa, N. Vittorio, B. D. Wandelt, I. K. Wehus, M. White, S. D. M. White, A. Zacchei, and A. Zonca, Planck 2018 results. VI. Cosmological parameters, A&A **641**, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [48] E. S. Phinney, A Practical Theorem on Gravitational Wave Backgrounds, ArXiv Astrophysics e-prints (2001), astro-ph/0108028.
- [49] R. W. Hellings and G. S. Downs, Upper limits on the isotropic gravitational radiation background from pulsar timing analysis, ApJ 265, L39 (1983).
- [50] R. Jinno and M. Takimoto, Gravitational waves from bubble collisions: An analytic derivation, Phys. Rev. D 95, 024009 (2017), arXiv:1605.01403 [astro-ph.CO].
- [51] M. Hindmarsh, S. J. Huber, K. Rummukainen, and D. J. Weir, Shape of the acoustic gravitational wave power spectrum from a first order phase transition, Phys. Rev. D 96, 103520 (2017), [Erratum: Phys.Rev.D 101, 089902 (2020)], arXiv:1704.05871 [astro-ph.CO].

- [52] M. Lewicki and V. Vaskonen, Gravitational waves from colliding vacuum bubbles in gauge theories, (2020), arXiv:2012.07826 [astro-ph.CO].
- [53] D. Cutting, E. G. Escartin, M. Hindmarsh, and D. J. Weir, Gravitational waves from vacuum first order phase transitions II: from thin to thick walls, Phys. Rev. D 103, 023531 (2021), arXiv:2005.13537 [astro-ph.CO].
- [54] C. Caprini et al., Science with the space-based interferometer eLISA. II: Gravitational waves from cosmological phase transitions, JCAP 04, 001, arXiv:1512.06239 [astro-ph.CO].
- [55] A. Roper Pol, S. Mandal, A. Brandenburg, T. Kahni-ashvili, and A. Kosowsky, Numerical simulations of gravitational waves from early-universe turbulence, Phys. Rev. D 102, 083512 (2020), arXiv:1903.08585 [astro-ph.CO].
- [56] A. Brandenburg, E. Clarke, Y. He, and T. Kahniashvili, Can we observe the QCD phase transition-generated gravitational waves through pulsar timing arrays?, Phys. Rev. D 104, 043513 (2021), arXiv:2102.12428 [astro-ph.CO].
- [57] P. Binetruy, A. Bohe, C. Caprini, and J.-F. Dufaux, Cosmological Backgrounds of Gravitational Waves and eLISA/NGO: Phase Transitions, Cosmic Strings and Other Sources, JCAP 06, 027, arXiv:1201.0983 [gr-qc].
- [58] H.-K. Guo, K. Sinha, D. Vagie, and G. White, Phase Transitions in an Expanding Universe: Stochastic Gravitational Waves in Standard and Non-Standard Histories, JCAP 01, 001, arXiv:2007.08537 [hep-ph].
- [59] J. Ellis, M. Lewicki, and J. M. No, Gravitational waves from first-order cosmological phase transitions: lifetime of the sound wave source, JCAP 07, 050, arXiv:2003.07360 [hep-ph].
- [60] D. J. Weir, Gravitational waves from a first order electroweak phase transition: a brief review, Phil. Trans. Roy. Soc. Lond. A 376, 20170126 (2018), arXiv:1705.01783 [hep-ph].
- [61] M. Hindmarsh and M. Hijazi, Gravitational waves from first order cosmological phase transitions in the Sound Shell Model, JCAP 12, 062, arXiv:1909.10040 [astroph.CO].
- [62] J. R. Espinosa, T. Konstandin, J. M. No, and G. Servant, Energy Budget of Cosmological First-order Phase Transitions, JCAP 06, 028, arXiv:1004.4187 [hep-ph].
- [63] A. Kosowsky, M. S. Turner, and R. Watkins, Gravitational radiation from colliding vacuum bubbles, Phys. Rev. D 45, 4514 (1992).
- [64] D. Cutting, M. Hindmarsh, and D. J. Weir, Gravitational waves from vacuum first-order phase transitions: from the envelope to the lattice, Phys. Rev. D 97, 123513 (2018), arXiv:1802.05712 [astro-ph.CO].
- [65] M. Lewicki and V. Vaskonen, Gravitational wave spectra from strongly supercooled phase transitions, Eur. Phys. J. C 80, 1003 (2020), arXiv:2007.04967 [astro-ph.CO].
- [66] G. Schwarz, Estimating the Dimension of a Model, Annals of Statistics 6, 461 (1978).
- [67] R. E. Kass and A. E. Raftery, Bayes factors, Journal of the American Statistical Association 90, 773 (1995).
- [68] D. Cutting, M. Hindmarsh, and D. J. Weir, Vorticity, kinetic energy, and suppressed gravitational wave production in strong first order phase transitions, Phys. Rev. Lett. 125, 021302 (2020), arXiv:1906.00480 [hep-ph].
- [69] Particle Data Group, P. A. Zyla, R. M. Barnett, J. Beringer, O. Dahl, D. A. Dwyer, D. E. Groom, C. J. Lin, K. S. Lugovsky, E. Pianori, D. J. Robinson, C. G.

Wohl, W. M. Yao, K. Agashe, G. Aielli, B. C. Allanach, C. Amsler, M. Antonelli, E. C. Aschenauer, D. M. Asner, H. Baer, S. Banerjee, L. Baudis, C. W. Bauer, J. J. Beatty, V. I. Belousov, S. Bethke, A. Bettini, O. Biebel, K. M. Black, E. Blucher, O. Buchmuller, V. Burkert, M. A. Bychkov, R. N. Cahn, M. Carena, A. Ceccucci, A. Cerri, D. Chakraborty, R. S. Chivukula, G. Cowan, G. D'Ambrosio, T. Damour, D. de Florian, A. de Gouvêa, T. DeGrand, P. de Jong, G. Dissertori, B. A. Dobrescu, M. D'Onofrio, M. Doser, M. Drees, H. K. Dreiner, P. Eerola, U. Egede, S. Eidelman, J. Ellis, J. Erler, V. V. Ezhela, W. Fetscher, B. D. Fields, B. Foster, A. Freitas, H. Gallagher, L. Garren, H. J. Gerber, G. Gerbier, T. Gershon, Y. Gershtein, T. Gherghetta, A. A. Godizov, M. C. Gonzalez-Garcia, M. Goodman, C. Grab, A. V. Gritsan, C. Grojean, M. Grünewald, A. Gurtu, T. Gutsche, H. E. Haber, C. Hanhart, S. Hashimoto, Y. Hayato, A. Hebecker, S. Heinemeyer, B. Heltsley, J. J. Hernández-Rey, K. Hikasa, J. Hisano, A. Höcker, J. Holder, A. Holtkamp, J. Huston, T. Hyodo, K. F. Johnson, M. Kado, M. Karliner, U. F. Katz, M. Kenzie, V. A. Khoze, S. R. Klein, E. Klempt, R. V. Kowalewski, F. Krauss, M. Kreps, B. Krusche, Y. Kwon, O. Lahav, J. Laiho, L. P. Lellouch, J. Lesgourgues, A. R. Liddle, Z. Ligeti, C. Lippmann, T. M. Liss, L. Littenberg, C. Lourengo, S. B. Lugovsky, A. Lusiani, Y. Makida, F. Maltoni, T. Mannel, A. V. Manohar, W. J. Marciano, A. Masoni, J. Matthews, U. G. Meißner, M. Mikhasenko, D. J. Miller, D. Milstead, R. E. Mitchell, K. Mönig, P. Molaro, F. Moortgat, M. Moskovic, K. Nakamura, M. Narain, P. Nason, S. Navas, M. Neubert, P. Nevski, Y. Nir, K. A. Olive, C. Patrignani, J. A. Peacock, S. T. Petcov, V. A. Petrov, A. Pich, A. Piepke, A. Pomarol, S. Profumo, A. Quadt, K. Rabbertz, J. Rademacker, G. Raffelt, H. Ramani, M. Ramsev-Musolf, B. N. Ratcliff, P. Richardson, A. Ringwald, S. Roesler, S. Rolli, A. Romaniouk, L. J. Rosenberg, J. L. Rosner, G. Rybka, M. Ryskin, R. A. Ryutin, Y. Sakai, G. P. Salam, S. Sarkar, F. Sauli, O. Schneider, K. Scholberg, A. J. Schwartz, J. Schwiening, D. Scott, V. Sharma, S. R. Sharpe, T. Shutt, M. Silari, T. Sjöstrand, P. Skands,

- T. Skwarnicki, G. F. Smoot, A. Soffer, M. S. Sozzi, S. Spanier, C. Spiering, A. Stahl, S. L. Stone, Y. Sumino, T. Sumiyoshi, M. J. Syphers, F. Takahashi, M. Tanabashi, J. Tanaka, M. Taševský, K. Terashi, J. Terning, U. Thoma, R. S. Thorne, L. Tiator, M. Titov, N. P. Tkachenko, D. R. Tovey, K. Trabelsi, P. Urquijo, G. Valencia, R. Van de Water, N. Varelas, G. Venanzoni, L. Verde, M. G. Vincter, P. Vogel, W. Vogelsang, A. Vogt, V. Vorobyev, S. P. Wakely, W. Walkowiak, C. W. Walter, D. Wands, M. O. Wascko, D. H. Weinberg, E. J. Weinberg, M. White, L. R. Wiencke, S. Willocq, C. L. Woody, R. L. Workman, M. Yokoyama, R. Yoshida, G. Zanderighi, G. P. Zeller, O. V. Zenin, R. Y. Zhu, S. L. Zhu, F. Zimmermann, J. Anderson, T. Basaglia, V. S. Lugovsky, P. Schaffner, and W. Zheng, Review of Particle Physics, Progress of Theoretical and Experimental Physics 2020, 083C01 (2020).
- [70] J. A. Ellis, M. Vallisneri, S. R. Taylor, and P. T. Baker, ENTERPRISE: Enhanced Numerical Toolbox Enabling a Robust PulsaR Inference SuitE (2019), ascl:1912.015.
- [71] S. R. Taylor, P. T. Baker, J. S. Hazboun, J. J. Simon, and S. J. Vigeland, enterprise extensions (2018).
- [72] J. Hazboun, J. Romano, and T. Smith, Hasasia: A Python package for Pulsar Timing Array Sensitivity Curves, The Journal of Open Source Software 4, 1775 (2019).
- [73] M. Vallisneri, libstempo: Python wrapper for Tempo2 (2020), ascl:2002.017.
- [74] J. D. Hunter, Matplotlib: A 2D Graphics Environment, Computing in Science and Engineering 9, 90 (2007).
- [75] J. Ellis and R. van Haasteren, jellis18/ptmcmcsampler: Official release (2017).
- [76] D. Nice, P. Demorest, I. Stairs, R. Manchester, J. Taylor, W. Peters, J. Weisberg, A. Irwin, N. Wex, and Y. Huang, Tempo: Pulsar timing data analysis (2015), ascl:1509.002.
- [77] G. Hobbs and R. Edwards, Tempo2: Pulsar Timing Package (2012), ascl:1210.015.
- [78] J. Luo, S. Ransom, P. Demorest, R. van Haasteren, P. Ray, K. Stovall, M. Bachetti, A. Archibald, M. Kerr, J. Colen, and F. Jenet, PINT: High-precision pulsar timing analysis package (2019), ascl:1902.007.